

THE COMMISSIONER OF PATENTS AND TRADEMARKS

WASHINGTON D.C. 20231

Case Docket No. 00/20989

Sir:

Transmitted herewith for filing is the Patent Application of

Inventor : Yair FELD; Lior EPSTEIN; Shimon MAROM; Meira FRANK

FOR : NUCLEIC ACID CONSTRUCTS AND CELLS, AND METHOD UTILIZING SAME FOR MODIFYING THE ELECTROPHYSIOLOGICAL FUNCTION OF EXCITABLE TISSUES

Enclosed are:

- ☒ 8 sheets of drawings
- ☒ An Assignment to: Technion Research and Development Foundation Ltd.
- ☒ Applicant is entitled to Small Entity Status under 37 CFR 1.9 and 37 CFR 1.27

THE FILING FEE HAS BEEN CALCULATED AS SHOWN BELOW:

FOR:	Column 1 No. FILED	Column 2 No. EXTRA	SMALL ENTITY		OTHER THAN A SMALL ENTITY	
			RATE	FEE	RATE	FEE
Basic Fee				\$ 355		\$ 710
Total Claims	37 -20=	17	17 x \$ 9	\$ 153	x \$18	\$
Independent Claims	4 - 3=	1	1 x \$40	\$ 40	x \$78	\$
<input checked="" type="checkbox"/> Recordal of Assignment \$40			+\$40	\$ 40	+\$40	\$
(If the difference in Column 1 is less than "0", enter "0" in Col. 2)			707AL	\$ 588	707AL	\$

☒ Please charge my deposit account No. 50-1407 in the amount of \$ 588.-.

A duplicate copy of this sheet is enclosed.

☒ The Commissioner is hereby authorized to charge payment of the following fees associated with this communication or credit any overpayment to Deposit Account No 50-1407. A duplicate copy of this sheet is enclosed.

☒ Any additional filing fees required under 37 CFR 1.16.

☒ Any patent application processing fees under 37 CFR 1.17.

☒ The Commissioner is hereby authorized to charge payment of the following fees during the pendency of this application or credit any overpayment to deposit Account No. 50-1407. A duplicate copy of this sheet is enclosed.

☒ Any patent application processing fees under 37 CFR 1.17

☐ The issue fee set in 36 CFR 1.18 at or before mailing of the Notice of Allowance, pursuant to 37 CFR 311(b).

☒ Any filing fees under 37 CFR 1.16 for presentation of extra claims.

Respectfully,

Sol Sheinbein

Sol Sheinbein

Registration No. 25,457

10/20/00

jc685 U.S. PTO

jc542 U.S. PTO
09/691889

10/20/00

09691889 102000

APPLICATION FOR PATENT

Inventors: Yair Feld, Lior Gepstein, Shimon Marom, Meira Frank

Title:

NUCLEIC ACID CONSTRUCTS AND CELLS, AND
METHODS UTILIZING SAME FOR MODIFYING THE
ELECTROPHYSIOLOGICAL FUNCTION OF EXCITABLE
TISSUES

FIELD AND BACKGROUND OF THE INVENTION

The present invention relates to nucleic acid constructs and cells, and further to methods utilizing same for modifying the electrophysiological function of excitable tissues. More particularly, embodiments of the present invention relate to the use of cells having gap junctions and ion channels or transporters for modifying the electrophysiological function of excitable tissues.

The biological cell membrane, the interface between the cell and its environment, is a complex biochemical entity one of whose major involvement is the directed transport of specific substances. A related major involvement of the cell membrane is the maintenance of chemical gradients, particularly electrochemical gradients, across this interface.

These gradients are of great functional significance (e.g., in the production of action potentials in nerve and muscle cells).

Ion channels are macromolecular protein pores, which span the cell membrane lipid bilayer. While approximately 30% of the energy expended in cells goes to maintain the ionic gradient across the cell membrane, it is the ion channel that dissipates this stored energy, much as a switch releases the electrical energy of a battery.

Ion channels are efficient compared to enzymes; small conformational changes gate a single channel between “closed” and “open” states, allowing up to 10^7 ions to flow in one second, amounting to approximately 10^{-12} Amperes of highly selected ions flow during the channel opening. Since they are efficient, the number of ion channels per cell is relatively low; a few thousand channels of a given subtype/cell are usually sufficient to perform their task while orders of magnitude higher numbers of receptors or enzymes are required to carry out their tasks.

Ion channels are usually classified by the type of ion they selectively pass (sodium, potassium, calcium, or chloride) although some are indiscriminate. Different ion channels are activated (or gated) by either extracellular ligands, transmembrane voltage, or intracellular second messengers.

Ion channel conductance

Conductance quantifies the ease with which ions flow through a material and is expressed in units of charge/sec/volt. Single channel conductance, g , as distinguished from the membrane conductance (G) of the entire population of channels, is defined as the ratio of single channel current amplitude (i) to the electromotive force, or voltage (V):

$$g=i/V$$

The direction of ion movement through channels is governed by electrical and chemical concentration gradients. Entropy dictates that ions will flow passively through ion channels down a chemical gradient. Electrically charged ions will also move in an electrical field, just as ions in solution flow to one of the poles of a battery connected to the solution. The point at which the chemical driving force is just balanced by the electrical driving force is called the Nernst equilibrium (or reversal) potential. Above or below this point, a particular ion species will flow in the direction of the dominant force. The net electrical flow across a cell membrane is predictable given the concentrations of ions, the number, conductances, and selectivities of the channels, and their gating properties.

The modern method of deciphering ion channel function is by using patch clamp technology. In the patch clamp technique, a small polished electrode is pressed against the plasma membrane. For unknown

reasons, the affinity between glass and cellular membrane is incredibly high; very few ions leak through this tight seal. In essence, the electrode isolates and captures all ions flowing through the 1-3 square microns of the cell membrane defined by the circular border of the glass pipette. The result is that the ionic current passing through a single ion channel can be collected and measured. The current through the attached patch (cell-attached), a detached patch (inside-out or outside-out), or the whole cell can be measured.

Ion channel building blocks

Since ion channel function is easily measured in real time, most ion channels were cloned using the South African clawed toad (*Xenopus laevis*) oocyte. These oocytes are large enough to inject with exogenous mRNA and are capable of synthesizing the resulting foreign proteins. In expression cloning, in vitro transcripts (mRNA) from a cDNA library derived from a source of tissue/cell known to be rich in a particular current are injected into individual oocytes. The proteins encoded by this library are allowed several days to be translated and processed before the oocyte currents are measured by voltage clamp techniques. The cDNA library (with ~ 1 million unique clones) is serially subdivided until injected messenger RNA from a single cDNA clone is isolated that confers novel ion channel activity. Moreover, mutant cDNA clones with engineered alterations in the protein's primary structure can be expressed and the ion

channel properties studied in order to determine regions of the protein critical for channel activation, inactivation, ion permeation, or drug interaction.

The building blocks for most channel proteins are individual polypeptide subunits or domains of subunits each containing six hydrophobic transmembrane regions labeled S1 through S6. The Na⁺ and Ca²⁺ channel pores are single (a) subunits in which 4 repeats of the six transmembrane spanning domain surround the pore. Voltage-gated K⁺ channels (Kv; nomenclature refers to K channel, voltage-dependent) are encoded by a tetramer of separate six-transmembrane spanning motifs. Coassembly of the linked domains form the central pore and confer the basic gating and permeation properties characteristic of the channel type. The peptide chain (H5 or P loop) juxtaposed between the membrane spanning segments S5 and S6 project into and line the water-filled channel pore. Mutations in this region alter the channel's permeation properties. S4 is probably the major channel voltage sensor since it contains a cluster of positively charged amino acids (lysines and arginines). Voltage-dependent channel inactivation is mediated by a tethered amino terminal blocking particle (called the ball and chain mechanism) which swings in to occlude the permeation pathway (inactivation). Amino acids in the S6 transmembrane segment participate in another inactivation pathway named C-type inactivation.

000201" 68816960

The most recently discovered family of channel proteins are the inward rectifier K⁺-selective channels (Kir; K channel, inward rectifier). These channels determine the resting membrane potential in most cells because they are open at rest. Kir channel topography is similar to the Kv channel class but the subunits lack the S1 to S4 segments present in Kv channels. The two transmembrane spanning domain surrounding the conserved H5 pore domain is deceptively simple; heteromultimeric channel formation, direct G protein gating, and interactions with other proteins by some Kir subtypes considerably increases the complex behavior of this channel class.

Ion transporters:

Yet another class of molecules which participate in ion transport across cellular membranes are the ion transporters. Ion transporters are integral membrane proteins capable of pumping one ion out of the cell while pumping another ion into the cell.

In, for example, Na/K ion transporters, the Na⁺, K⁺ pump activity is the result of an integral membrane protein called the Na⁺, K⁺ -ATPase. The Na⁺, K⁺ -ATPase consists of a “catalytic” α-subunit of about 100,000 daltons and a glycoprotein β-subunit of about 50,000 daltons. When operating near its capacity for ion transport, the Na⁺, K⁺ -ATPase transport three sodium ions out of the cell and transport two potassium ions into the cell for each ATP hydrolyzed. The cyclic phosphorylation

and dephosphorylation of the protein causes it to alternate between two conformations, E1 and E2. In E1 the ion-binding sites of the protein have high affinity for Na^+ and face the cytoplasm. In the E2 conformation the ion-binding sites favor the binding of K^+ and face the extracellular fluid.

5 Examples of other ion transporters include the Na/Ca exchange system which participates in regulation of intracellular Ca^{++} ; the Na/H exchange system which function in concert with a Cl/HCO_3^- exchange system to regulate intracellular pH; and the Na-K-Cl exchange system which contributes to smooth muscle function and which is regulated by a
10 number of vasoactive agents.

Excitable tissues

Myocardium: Myocardial contraction depends on the opening and closing of a complex series of ion channels in myocardial cell membranes.

The most prominent of these channels are the K^+ , Ca^{++} and Na^+ ion
15 channels.

The number of K^+ ions is greater inside a resting myocardial cell than outside. But the number of Na^+ ions is greater outside. When a myocardial cell beats, sodium channels open allowing a rapid, transient in-rush of Na^+ ions, then close within about two one-thousandth's ($2/1000$)
20 of a second. This depolarizes the membrane with the positive ions moving in. Then there is then a slower, but prolonged ($1/2$ second), release of potassium to the outside of the cell which repolarizes the cell membrane.

Although myocardial contraction is more complex and involves other ions and channels, the end result of this depolarization-repolarization is that the contractile filaments in the cell engage, and the cell contracts.

5 **Nerve cells:** Signal propagation through neuronal cells is also governed by ion influx/efflux through nerve cell membranes. In nerve cells, Na^+ , Ca^{++} and K^+ channels participate in the generation and propagation of a nerve signal.

10 **Glandular tissue:** Secretion of glandular factors, such as hormones is in some cases effected by the excitation of secreting cells or tissues. For example, in the pancreas, T-type calcium channels along with cell-to-cell gap junctions participate in secretion of insulin.

Since ion channels participate in numerous physiological processes, damage to cells and/or channels of excitable tissues can be a
15 cause for numerous disorders.

For example, heart conditions, such as reentrant arrhythmia, are brought about by the damage or death of myocardial cells, which can no longer support normal electrophysiological function. Secretion of factors from glandular tissue, such as insulin is also effected by damage to
20 excitable cells forming this tissue, while nerve cell changes, as for instance in disorders such as, epilepsy severely effects nerve signal function.

The present invention provides a novel approach for modifying the electrophysiological property and thus the electrophysiological function of excitable tissues.

This novel approach, which according to one embodiment of the present invention utilizes cellular implants, can be utilized for restoring normal electrophysiological function to damaged tissues such as heart, nerve or glandular tissues.

SUMMARY OF THE INVENTION

According to one aspect of the present invention there is provided a nucleic acid construct comprising: (a) a first polynucleotide region encoding at least one first polypeptide capable of forming a functional ion channel or transporter when expressed within a cell; and (b) a second polynucleotide region encoding at least one second polypeptide capable of forming a functional gap junction when expressed within the cell.

According to further features in preferred embodiments of the invention described below, the nucleic acid construct further comprising at least one promoter being for directing the transcription of the first polynucleotide and the second polynucleotide.

According to still further features in the described preferred embodiments the at least one promoter is functional in mammalian cells.

According to still further features in the described preferred embodiments the at least one promoter is selected from the group consisting of a constitutive promoter, a tissue specific promoter, an inducible promoter and a developmentally regulated promoter.

According to still further features in the described preferred embodiments the first polynucleotide region and the second polynucleotide region are transcriptionally fused via an IRES sequence.

According to still further features in the described preferred embodiments the at least one first polypeptide and the at least one second polypeptide are translationally fused via at least one protease recognition site.

According to still further features in the described preferred embodiments the at least one promoter includes two promoters, a first promoter for directing the transcription of the first polynucleotide and a second promoter for directing the transcription of the second polynucleotide.

According to another aspect of the present invention there is provided a nucleic acid construct system comprising: (a) a first nucleic acid construct including a first polynucleotide region encoding at least one first polypeptide capable of forming a functional ion channel or transporter when expressed within a cell; and (b) a second nucleic acid construct including a second polynucleotide region encoding at least one

second polypeptide capable of forming a functional gap junction when expressed within the cell.

According to still further features in the described preferred embodiments the first nucleic acid construct further includes a first
5 promoter being for directing the transcription of the first polynucleotide and further wherein the second nucleic acid construct further includes a second promoter being for directing the transcription of the second polynucleotide.

According to still further features in the described preferred embodiments each of the first and the second promoters is functional in mammalian cells.

According to still further features in the described preferred
10 embodiments each of first and the second promoters is independently selected from the group consisting of a constitutive promoter, a tissue specific promoter, an inducible promoter and a developmentally regulated promoter.

According to still further features in the described preferred embodiments there is provided a cell, cell culture or tissue explant transformed with the nucleic acid constructs described above.

According to still further features in the described preferred embodiments the cell is selected from the group consisting of a fibroblast, a myoblast, an astroglial cell and an endothelial cell.

000201" 68876560

According to still further features in the described preferred embodiments the tissue explant is an organ tissue explant.

According to still further features in the described preferred embodiments there is provided a pharmaceutical composition comprising, as an active ingredient, the nucleic acid constructs described above.

According to still further features in the described preferred embodiments the ion channel is selected from the group consisting of a sodium ion channel, a potassium ion channel, a calcium ion channel and a chloride ion channel.

According to still further features in the described preferred embodiments the at least one first polypeptide is selected from the group consisting of a K channel polypeptide, a Na channel polypeptide, a Ca channel polypeptide, a Cl channel polypeptide, a Na/K transporter polypeptide, a Na/Ca transporter polypeptide, a Na/H transporter polypeptide and a Cl/ HCO₃ transporter polypeptide.

According to still further features in the described preferred embodiments the at least one second polypeptide is selected from the group consisting of connexin43, connexin45 and connexin26.

According to still another aspect of the present invention there is provided a method of modifying the electrophysiological function of an excitable tissue region of an individual, the method comprising the step of implanting cells into the excitable tissue region, each implanted cell being:

(a) capable of forming gap junctions with at least one cell of the excitable tissue region; and (b) capable of forming a functional ion channel or transporter; the functional ion channel or transporter being capable of modifying the electrophysiological function of the excitable tissue region.

According to still further features in the described preferred embodiments the ion channel is selected from the group consisting of a sodium ion channel, a potassium ion channel, a calcium ion channel and a chloride ion channel.

According to still further features in the described preferred embodiments each implanted cell is transfected, prior to, or following implantation, with an exogenous polynucleotide expressing at least one polypeptide capable of forming the functional ion channel or transporter.

According to still further features in the described preferred embodiments each implanted cell is transformed, prior to, or following implantation, with an exogenous polynucleotide expressing at least one polypeptide capable of forming the gap junctions.

According to still further features in the described preferred embodiments expression of the at least one polypeptide from the exogenous polynucleotide is regulatable by an endogenous or an exogenous factor.

According to still further features in the described preferred embodiments an ion permeability of the functional ion channels is regulatable by an endogenous or an exogenous factor.

According to still further features in the described preferred embodiments the method further comprising the step of regulating the permeability of the functional ion channels, or the activity of the transporter to thereby regulate the electrophysiological function of the excitable tissue region.

According to still further features in the described preferred embodiments the step of regulating the permeability is affected by providing the exogenous factor to the excitable tissue region.

According to still further features in the described preferred embodiments each implanted cell is capable of forming the functional ion channel or transporter following induction.

According to still further features in the described preferred embodiments the excitable tissue region forms a part of an organ selected from the group consisting of a heart, a pancreas, a kidney, a brain and a liver.

According to still further features in the described preferred embodiments the method is utilized for regulating cardiac arrhythmia.

According to still further features in the described preferred embodiments the method is utilized for regulating secretion of

endogenous factors from an organ including the excitable tissue region of the individual.

According to still further features in the described preferred embodiments the method is utilized for regulating neuronal discharge.

According to an additional aspect of the present invention there is provided a method of modifying the electrophysiological function of an excitable tissue region of an individual, the method comprising the step of expressing an exogenous polypeptide in at least a portion of cells forming a part of, or being in contact with, the excitable tissue region, the exogenous polypeptide being capable of forming functional ion channels or transporters within the at least a portion of the cells to thereby modify the electrophysiological function of the excitable tissue region.

According to still further features in the described preferred embodiments the method further comprising the step of expressing a second exogenous polypeptide in the at least a portion of the cells, the second exogenous polypeptide being capable of forming functional pap
 5 junctions within the at least a portion of the cells.

The present invention successfully addresses the shortcomings of the presently known configurations by providing a novel approach for modifying the electrophysiological function of excitable tissues.

BRIEF DESCRIPTION OF THE DRAWINGS

The invention is herein described, by way of example only, with reference to the accompanying drawings. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of the preferred embodiments of the present invention only, and are presented in the cause of providing what is believed to be the most useful and readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show structural details of the invention in more detail than is necessary for a fundamental understanding of the invention, the description taken with the drawings making apparent to those skilled in the art how the several forms of the invention may be embodied in practice.

In the drawings:

FIGs. 1a-e illustrate results from experiments performed on large random cortical networks cultured on substrate-embedded multi-electrode arrays (MEA). Figure 1a-b - an image illustrating four out of sixty electrodes and the somata of numerous neurons growing on the surface (1a) and a magnification of a region thereof (1b). In the magnified image (1b) the richness of the connective (axo-dendritic) network is evident.

Scale bar: 30 μm . Figure 1c - exemplifies an action potential recorded from one electrode. The two parallel lines represent $\pm 8\text{RMS}$ units for this particular electrode. Network response to focal stimulation. Figures 1d-e illustrate a reverberating response of the network to focal stimuli. A typical stimulus pulse lasts 420 μSec , and its amplitude is 50 μA . The traces were recorded simultaneously from different electrodes. Note the reverberating response to a stimulus (enlarged in Figure 1e) which lasts 100 milliseconds or more. Figure 1f is a graph illustrating the connectivity in cultured networks. The average number (four networks) of significantly occurring activity pairs formed between ten randomly chosen active ($>0.2\text{Hz}$ of spontaneous activity) electrodes. This number, normalized to the maximal number of possible activity pairs, is depicted as *fraction connected*, and shown to decrease as a function of within-pair time delay (δ). Inset: Given an $A \rightarrow B$ activity pair, the forecasting of B by A , which is the strength of the functional connectivity between the two, is given in terms of a correlation coefficient. This correlation is calculated from the number of times that the given pair appears within 1 hour, divided by the number of occurrences of A OR B . The average ($n=4$) functional connectivity strength as a function of δ is shown.

FIG. 2a illustrates epileptic activity recorded from MEA in a mature (3 weeks in vitro) cultured cortical network. The network is prepared and recorded from as explained in Figure 1. The recorded spontaneously

bursting synchronous activity throughout the network is a characteristic feature of epileptic-like activity in networks of neurons.

FIG. 2b illustrates an expanded time scale of the activity marked by the red box (left side) revealing a complex structure of a single burst.

FIG. 3 illustrates the dose response to charybdotoxin of the Kv1.3 potassium channel.

FIG. 4a illustrates a diffused fibroblast seeding pattern on multi electrode array (fibroblasts in red).

FIG. 4b illustrates a clustered fibroblast seeding pattern on multi electrode array.

FIG. 5 illustrates the CTX frequency response of cultured cardiomyocytes (square), cardiomyocytes co-cultured with fibroblasts NIH 3T3 (dot) and cardiomyocytes co-cultured with fibroblasts (NIH 3T3) transfected with voltage gated potassium channel Kv1.3 coding sequence (triangle). c-no. of cultures; n-no. of measurements (without fibroblast - c=8; n=16, NIH 3T3 - c=6; n=47, Kv1.3 -c=6; n=43, error bar – standard error).

FIG. 6a is a fluorescent image of a cardiomyocytes co-cultured with fibroblasts transfected with Kv1.3 channel coding sequences and labeled with Fast DiO (MAE cluster seeding pattern). The blue dot marks electrode 28 and the red dot marks electrode 53.

FIGs. 6b-c represent a two second recording of synchronous extracellular activity prior to seeding of the fibroblasts described in Figure 6a. Figure 6b- recording from electrode 28; Figure 6c- recording from electrode 53;

5 FIGs. 6d-e represent a two second recording of uncoupled extracellular activity following seeding of the fibroblasts described in Figure 6a and prior to treatment with CTX. Figure 6d- recording from electrode 28; Figure 6e- recording from electrode 53;

10 FIG. 6f-g represent a two second recording of extracellular activity following seeding of the fibroblasts described in Figure 6a and treatment with CTX 100 nM which reverses uncoupling effect. Figure 6f- recording from electrode 28; Figure 6g- recording from electrode 53;

15 FIG. 7a is a fluorescent image of cardiomyocytes co-cultured with fibroblasts transfected with Kv1.3 channel coding sequences and labeled with Fast DiO on a MEA (cluster seeding pattern).

FIG. 7b illustrates an activation map constructed prior to seeding of the fibroblasts described in Figure 7a.

20 FIG. 7c is an activation map constructed five days following seeding of the fibroblasts described in Figure 7a and prior to treatment with CTX illustrating the appearance of a conduction block.

FIG. 7d is an activation map constructed five days following seeding of the fibroblasts described in Figure 7a and following treatment

with CTX 10 nM illustrating the reversal of the conduction block.

FIG. 8 illustrates the conduction velocity change throughout an experiment with seeded fibroblasts (in blue – myocytes with fibroblasts transfected with Kv1.3, in brown – myocytes with fibroblast without Transfection (control 1), in yellow – myocytes without fibroblast (control 2)) ; fibroblasts where seeded following measurements at day 1; Kv1.3 c=4, NIH 3T3 c=3, without fib c=1 error bar – standard error.

FIG. 9 illustrates the amplitude change throughout the experiment illustrated in Figure 8; fibroblasts where seeded following measurements at day 1; Kv1.3 c=3, NIH 3T3 c=3 error bar – standard error.

FIG. 10 illustrates the development of a conduction block in MEA seeded fibroblasts following measurement at day 0. A substantial increase in the conduction block factor was recorded from the culture including the fibroblast transfected with potassium channels (Kv1.3) (pink, n=5), while in the non-transfected fibroblast culture a decrease in the conduction block factor was recorded (blue, n=6).

FIG.11 illustrates the effects of Charybdotoxin (specific blocker of potassium channel Kv1.3) on conduction blocks. In co-cultures including fibroblasts transfected with Kv1.3, application of Charybdotoxin substantially decreased the conduction block factor (pink, n=10), while in co-cultures including non-transfected fibroblasts, a minimal response was recorded (blue, n=9).

FIG.12a is a fluorescent image of MEA cultured cardiomyocytes and fibroblasts labeled with Fast DiO (cluster seeding pattern).

FIG. 12b illustrates an activation map constructed prior to seeding of the fibroblasts described in Figure 12a.

5 FIG. 12c illustrates an activation map constructed five days following seeding of the fibroblasts described in Figure 12a and prior to treatment with CTX, no conduction block is apparent.

FIG. 12d illustrates an activation map constructed five days following seeding of the fibroblasts described in Figure 12a and following
10 treatment with CTX (10 nM); no appreciable change from the activation map of Figure 12c is evident.

15 DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention is of nucleic acid constructs and cells, and of methods utilizing same for modifying the electrophysiological function of excitable tissues. Specifically, the present invention can be used to restore normal electrophysiological function to cells or tissues of, for
20 example, damaged myocardium, neurons and secretory glands.

The principles and operation of the present invention may be better understood with reference to the accompanying descriptions.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not limited in its application to the details of construction and the arrangement of the components set forth in the following description. The invention is capable of other
5 embodiments or of being practiced or carried out in various ways. Also, it is to be understood that the phraseology and terminology employed herein is for the purpose of description and should not be regarded as limiting.

Since electrophysiological function of excitable tissues is governed
10 by the quantity and type of ion channels present in the membrane of cells forming the excitable tissue, as well as the presence of gap junctions networking these cells, the present inventors propose that the electrophysiological function of any excitable tissue region can be modified by either expressing ion channel/transporter polypeptide(s)
15 and/or gap junction polypeptide(s) within cells forming a part of, or being in contact with, the excitable tissue region or by implanting cells which possess ion channels/transporters and gap junctions within excitable tissues.

As used herein, the phrase "excitable tissue" refers to tissue which
20 is composed, at least in part, of cells which respond to, or propagate, an electrochemical change. Examples include muscle tissue, neuronal tissue and glandular tissue.

According to the present invention, the introduction of new channels or channel producing cells into an excitable tissue, as well as the regulation of channel formation or permeability via endogenous or exogenous factors, can be utilized to control the electrophysiological function of excitable tissue to thereby treat various disorders associated with such tissues.

Thus, according to one aspect of the present invention, there is provided a nucleic acid construct including a first polynucleotide region encoding at least one first polypeptide which is capable of forming a functional ion channel or transporter when expressed within a cell, and a second polynucleotide region encoding at least one second polypeptide capable of forming a functional gap junction when expressed within the cell.

According to a preferred embodiment of the present invention, the first polynucleotide region encodes an ion channel forming polypeptide or polypeptides, such as, but not limited to, a Ca, K, Na or Cl ion channel forming polypeptide(s). For example, the first polynucleotide region can include the sequence set forth by nucleotides 179-6121 of Genbank Accession number AB027567, which when expressed within the cell produces a Na channel.

Additional examples of sequences which can be utilized by the present invention for forming a functional ion channel, when expressed

within the cell, are listed according to their GenBank accession numbers in Tables 1-3 of the Example section which follows.

The first polynucleotide region can also encode any modified polypeptide (e.g. mutated, chimeric etc') which is capable of forming functional ion channel in cells. Examples of mutated ion channel forming sequences are given in the Examples section which follows.

It will be appreciated that ion transporters such as Na/K, Na/Ca or Cl/HCO₃ exchange systems (ATPases) can also be utilized by the present invention. Since such transporters are typically slower than channels in transporting ions across cell membranes, their use is limited to cases where rapid influx or outflux of ions is not required.

According to another preferred embodiment of the present invention, the gap junction forming polypeptide encoded by the second polynucleotide region is Connexin43 or 45, other connexin types which can be utilized by the present invention are described in the Examples section which follows.

The nucleic acid construct according to this aspect of the present invention also includes at least one promoter sequence for driving the transcription of the first and second polynucleotide regions. Preferably, the nucleic acid construct includes two promoters each driving transcription of a specific polynucleotide region. Alternatively, a single promoter sequence can transcribe both polynucleotide regions as a

polycistronic message. Such a polycistronic message can include an internal ribosome entry site (IRES) between the first and second polynucleotide regions so as to enable the translation of the downstream polynucleotide region. Alternatively, the first and second polynucleotide regions of the polycistronic message can be translationally fused via a protease recognition site, such that a polypeptide translated from this message is cleaved into the first and second polypeptides described above.

It will be appreciated that although expressing both polynucleotide regions from a single construct is advantageous in some respects, each of the polynucleotide regions can alternatively be provided on a separate construct.

Thus, according to another aspect of the present invention there is provided a nucleic acid construct system which includes a first nucleic acid construct including a first polynucleotide region encoding at least one first polypeptide capable of forming a functional ion channel or transporter when expressed within a cell and a second nucleic acid construct including a second polynucleotide region encoding at least one second polypeptide capable of forming a functional gap junction when expressed within the cell.

The nucleic acid constructs of the present invention are utilized to transform cells, preferably mammalian cells, either in-vivo or ex-vivo.

As such the promoters utilized by these construct are mammalian functional promoters which are either constitutive, tissue specific, inducible or growth regulatable depending on the cell type and application.

The nucleic acid constructs described hereinabove are preferably constructed using commercially available mammalian expression vectors or derivatives thereof. Examples of suitable vectors include, but are not limited to, pcDNA3, pcDNA3.1(+/-), pZeoSV2(+/-), pSecTag2, pDisplay, 5 pEF/myc/cyto, pCMV/myc/cyto, pCR3.1, which are available from Invitrogen, pCI which is available from Promega, pBK-RSV and pBK-CMV which are available from Stratagene, pTRES which is available from Clontech, and their derivatives and modificants.

Any of the promoter and/or regulatory sequences included in the mammalian expression vectors described above can be utilized to direct the transcription of the polynucleotide regions described above. However, since such vectors are readily amenable to sequence modifications via standard recombinant techniques, additional regulatory elements, promoter and/or selection markers can easily be incorporated therein if needed.

The nucleic acid constructs of the present invention can be introduced into a cell, population of cells, or tissue via any standard *in-vivo* or *ex-vivo* mammalian transformation method. Such methods include,

but are not limited to, direct DNA uptake techniques, and virus or liposome mediated transformation (for further detail see, for example, "Methods in Enzymology" Vol. 1-317, Academic Press).

The constructs according to the present invention can be administered to the individual *per se*, or in a pharmaceutical composition where it is mixed with suitable carriers or excipients.

Thus, according to another preferred embodiment of the present invention, the nucleic acid constructs according to the teachings of the present invention are included in a pharmaceutical composition which also includes a pharmaceutically acceptable carrier which serves for stabilizing and/or enhancing the accessibility or targeting of the constructs to target tissues.

As used herein a "pharmaceutical composition" refers to a preparation of one or more of the active ingredients described herein with other chemical components such as physiologically suitable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

Herein the term "active ingredient" refers to the preparation accountable for the biological effect, i.e. the nucleic acid constructs of the present invention.

Hereinafter, the phrases "physiologically acceptable carrier" and "pharmaceutically acceptable carrier" are interchangeably used to refer to

a carrier, such as, for example, a liposome, a virus, a micelle, or a protein, or a diluent which do not cause significant irritation to an organism and do not abrogate the biological activity and properties of the active ingredient. An adjuvant is included under these phrases.

5 Herein the term "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of an active ingredient. Examples, without limitation, of excipients, include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils and polyethylene glycols.

10 Techniques for formulation and administration of compositions may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition, which is incorporated herein by reference.

Suitable routes of administration are preferably local rather than systemic, for example, via injection of the preparation directly into the
15 excitable tissue region. For injection, the active ingredients of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological salt buffer.

Pharmaceutical compositions of the present invention may be
20 manufactured by processes well known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active ingredients into preparations which, can be used pharmaceutically.

Pharmaceutical compositions suitable for use in context of the present invention include compositions wherein the active ingredients are contained in an amount effective to achieve the intended purpose.

Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any preparation used in the methods of the invention, the therapeutically effective amount or dose can be estimated initially from *in vitro* and cell culture assays. For example, a dose can be formulated in animal models to achieve a desired concentration or titer of the active ingredient. Such information can be used to more accurately determine useful doses in humans.

Toxicity and therapeutic efficacy of the active ingredients described herein can be determined by standard pharmaceutical procedures *in vitro*, in cell cultures or experimental animals. The data obtained from these *in vitro* and cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage may vary depending

upon the dosage form employed. (See e.g., Fingl, *et al.*, 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1).

Direct administration of the nucleic acid constructs described hereinabove or of pharmaceutical compositions including such constructs
5 into cells forming a part of, or being in contact with, the excitable tissue region is preferably used in cases where the cells of the excitable tissue to be transformed are viable and functional.

In cases where cell damage or death defines a disorder of excitable tissue, the preferred mode of treatment is implantation of transformed or
10 non-transformed cells having ion channels/transporters and gap junctions.

Thus, according to another aspect of the present invention there is provided a method of modifying the electrophysiological function of an excitable tissue region of an individual. The method is effected by implanting cells into the excitable tissue region, wherein the implanted cells are each characterized by the ability to form gap junctions with at least one cell of the excitable tissue region and by the ability to form functional ion channels or transporters of one or more channel or transporter types.

Implantation of such cells can be effected by, for example, a syringe and needle adapted or fabricated for cell implantation, by a catheter drug delivery system (see for example, U.S. patent 6,102,887) or by standard

neurosurgical methods.

As mentioned above, the implanted cells can be cells expressing endogenous ion channel and/or gap junction polypeptides, or modified cells transformed with the nucleic acid constructs of the present invention. Preferably, the implanted cells are mammalian cells, such as for example, muscle, or fibers cells (see the Examples section for further detail).

In any case, the cells and ion channel selectivity and gating-regulation types are selected according to the application. For example, in application where rapid channel gating is crucial, an ion channel of regulatable gating is selected. Gating Regulated channels, and factors utilizable for regulating gating are described in the examples section hereinbelow.

In addition, regulation of ion channel/transporter polypeptide expression through, for example, induced promoter activity or the like can also be effected as an alternative or additive regulatory mechanism for controlling ion influx or outflux.

Thus, the present invention provides a novel approach for modifying the electrophysiological function of excitable tissues. As is further detailed in the Examples section which follows, the present invention can be utilized to restore enhance or suppress
5 electrophysiological function across a tissue region to thereby treat disorders caused by dysfunction in, or damage to, excitable tissues.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, cellular and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988); Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al.

(eds); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., eds. (1984); "Animal Cell Culture" Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); all of which are incorporated by reference as if fully set forth herein. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

15

EXAMPLE 1

Cardiac applications:

Cardiac arrhythmias are rhythm disturbances that result from alteration of the electrophysiological substrate of the heart. These arrhythmias include bradyarrhythmias (slow heart rate) which result from abnormalities in impulse formation or conduction and tachyarrhythmias (high heart rate) which result from abnormalities in the

electrophysiological substrate and which lead to the formation of tachycardia via abnormal foci firing at high rate or via formation of reentry circuits.

Cardiac arrhythmia often results from damage to the
 5 electrophysiological tissue substrate of the heart. By transplanting cells transfected with various ionic channels of specific and predetermined properties, the methods of the present invention enable one to modify the electrophysiological properties of heart tissue and thus repair such arrhythmias. Thus, the present invention can be used to either increase
 10 excitability to treat bradyarrhythmias or modify the electrophysiological substrate in order to suppress or prevent tachyarrhythmias.

Numerous cell types can be utilized to accomplish such a task, provided the cells possess functional gap junctions and functional ion channels.

15 Examples of suitable cell types include, but are not limited to, fibroblasts, skeletal myoblasts (satellite cells), endothelial cells and the like which can be of autogenic, allogenic, or xenogenic origin.

The cells transplanted generate specific structural and function interactions with the cardiomyocytes via the gap junction which can be
 20 either inherent to the transplanted cells or the product of overexpressed exogenes (listed in Table 1 below).

Table 1 - Sequences encoding polypeptide constituents of various ion channels

Ion	Channel type	GenBank Accession numbers	Potential application
K	Kv1.3	H18261	Reentrant arrhythmia, Atrial fibrillation, Ventricular and atrial tachycardia or heart failure
K	inward rectifier potassium channel TWIK-1 - human	S65566	Atrial fibrillation or heart failure
K	Delayed rectifier potassium channel - human	L28168 L33815 M26685	Atrial fibrillation or heart failure
K	Cardiac inward rectifier potassium channel - human	I38727	Atrial fibrillation or heart failure
K	VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KV1.4	M55514 A1631014 A1701825 A1694934 A1793138	Atrial fibrillation or heart failure
K	'voltage-gated potassium channel protein - human'	JC5275	Atrial fibrillation or heart failure
K	OKCNQ2"; potassium channel	AF033348	Atrial fibrillation or heart failure
K	'inwardly rectifying potassium channel, hippocampal	I38521	Reentrant arrhythmia, Atrial fibrillation, Ventricular and atrial tachycardia or heart failure
K	VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KQT-LIKE 3.KCNQ3.	AF033347 AF071491 AW205596 AW135705 AA019129 AA001392 H86059 H08544 R36327 T78692 A1125802 H08545 R49258	Atrial fibrillation or heart failure
Na	Sodium channel	AB027567	A-V block, Atrial fibrillation, Sick sinus syndrome
Na	Voltage gated "SCN11A"	AF188679	A-V block, Atrial fibrillation, Sick sinus syndrome
Na	'AMILORIDE-SENSITIVE BRAIN SODIUM CHANNEL BNAC1'	U57352 U50352 H12215 Z45660 R35720 R15377 AA457638 A1473139 H12216 A1825456 R49357 T16341 F04549 R42118	A-V block, Atrial fibrillation, Sick sinus syndrome
Na	hBNAC2"; product: "sodium channel 2	U78181 AL035862 AA442069 AI017398 AI620655 AI762424 Z40887 AI700050	A-V block, Atrial fibrillation, Sick sinus syndrome
Ca	T-type	AF134986	Heart failure
Ca	'VOLTAGE-DEPENDENT N-TYPE CALCIUM CHANNEL'	M94172 U76666 AA776162 T12610	Heart failure
Ca	"L-type calcium channel (HFCC)"; Human'	M92269 AA927640 AA443875 AA173146	Heart failure

000201" 588T6950

Ca	"CACNG4"; product: "calcium channel"	AF142625	Heart failure
Ca	"VOLTAGE-DEPENDENT L-TYPE CALCIUM CHANNEL,"	AJ224874 AJ006216	Heart failure
Ca	"voltage-dependent calcium channel"	M92301 W07059 T28094	Heart failure
Ca	L-type	M76558 AF055575 H29339 R25307 T27949 AA885750 AW029633 AI955764 AW008794 AA978315 AI914244 AI951788 AW008769 H29256 AI963788 AI537488 AA468565 AA523647 AI361691 R46658 AW139850 AI017959 AA701888 AA703120 AA877582	Heart failure
Cl	'probable chloride channel CIC-6 human'	S68428	
Cl	"CLCN3"; product: "chloride channel protein"	AF029346	
Cl	"CIC-2"; product: "chloride channel"	AF026004	
Cl	"clc4"; product: "chloride channel"	AB019432	

Table 1 Cont.

The coupling between the transplanted and host cells forms a single functional unit. Such functional coupling of the transplanted cells with the myocytic tissue allows modification of the various action potential phases of the myocytes.

Listed below are some of the action potential modifications, which can be effected using the methods of the present invention.

(i) Transplantation of fibroblasts having potassium channels (KV1-3,É) can be utilized to reduce automaticity; the effect may be reversed by specific antagonist (e.g., Charybdotoxin)

(ii) Transplantation of fibroblasts having potassium channels (KV1-3,É) can also be utilized for the creation of block which can be reversed with CTX.

(iii) Transplantation of fibroblasts having sodium channels can be utilized for the creation of rate dependent conduction block. Na channels will be inactivated at fast (abnormal) rates but permit conduction at slower (physiological) rates.

5 (iv) Transplantation of cells having various channels (for example the human ether-a-go-go-related gene, HERG) can be used to repress abnormal focal activity (due to triggered activity and unstable repolarization).

(v) Transplantation of cells having KV channels can be utilized
10 to regulate A-V node conduction (e.g., prolong refractoriness, or decrease conduction velocity).

(vi) Transplantation of cells having Na-channels or Na and K channels can be utilized to increase A-V node conduction.

(vii) Transplantation of cells having Na-channels can be utilized
15 to increase excitability by increasing spontaneous rate and conduction within the SA node (pacemaker).

Transplantation patterns:

The ability to transplant the cellular grafts of the present invention at predetermined myocardial sites may be of unique advantage since the
20 location of the transplantation site can be selected and optimized according to the specific mechanism of the arrhythmia treated.

For example, a local effect may decrease side effects which result from a more generalized effect, as occurs for example, during pharmacological treatments.

A focal transplantation pattern may be used to treat focal arrhythmia or change excitability at predetermined sites. Linear lesion transplantation may be utilized to generate conduction blocks for the treatment of specific reentrant arrhythmia while diffuse transplantation patterns may be utilized to modify the excitable properties of entire regions.

10 **Methods of transplantation:**

Several transplantation approaches can be utilized by the present invention. For example, an epicardial transplantation can be effected via surgical procedures, while an endocardial transplantation can be effected via catheters that are employed percutaneously and may be used to inject the cells endocardially. Alternatively, the cells may be injected into the coronary circulation.

Specific applications:

Atrial fibrillation (AF): In atrial fibrillation, the normal rhythmical contractions of the cardiac atria are replaced by rapid irregular twitchings of the muscular wall; the ventricles respond in an irregular and rapid manner to the dysrhythmic bombardment from the atria. The pathological

properties of AF can be modified using the teachings of the present invention via one of several possible approaches:

Cells transfected with specific ionic channel coding sequences, for example the voltage gated potassium channels (Kv1.3), can be transplanted into the A-V node. The modulating effect on the A-V node will slow the ventricular rate. This effect may be further modulated by dose-related changes resulting from the application of a blocking factor such as, for example, charybdotoxin.

The present invention also enables to treat AF by creating multiple line blocks (similar to the surgical maze procedure or the equivalent ablation procedure) in both atria. These blocks can be created by transplanting cells having K channels or rate dependent Na channels in the desired predetermined pattern.

The methods of the present invention cells may also be utilized to suppress pulmonary vein foci which often trigger AF, or to prevent their propagation to the atria by creating conduction blocks. By modifying the electrophysiological substrate of the atria the methods of the present invention can be used to increase cellular coupling and to increase and homogenize repolarization.

Atrial flutter and other Macroreentrant atrial arrhythmia: These arrhythmias result from macroreentrant wavefronts which can be treated

by transplanting the cells to create a block at a critical area (for example the tricuspid-IVC isthmus in typical flutter).

Atrial tachycardia: Paroxysmal tachycardia originating in an ectopic focus in the atrium can be treated by cells transplanted at the area of the ectopic foci which suppress the abnormal activity.

Ventricular and reentrant tachycardia: The methods of the present invention can also be utilized to treat paroxysmal tachycardia originating in an ectopic focus in the ventricle by transplanting cells at the area of the ectopic foci. In addition, reentrant tachycardia originating from a scar tissue, following myocardial infarction is also treatable via the methods of the present invention. In this case, cellular grafts can be used to modify (increase or decrease) the conduction properties of slow conduction pathways within the scar which are critical for initiation and sustainment of the reentrant arrhythmia.

A-V block: An impairment of the normal conduction between atria and ventricles can be treated by cellular graft which improve the excitability properties of the A-V node thus reversing the conduction block.

Sick-sinus syndrome: An abnormal function of the SA node (normal pace maker) which results in a slow heart rate or alternating slow-fast rates can be treated by cells transplanted in the SA Node area in order to increase the excitability of the SA Node, or by creating an alternative

pacemaker by transplanting cells with pace maker properties (combination of Na and K channels).

Heart failure: despite considerable advances in the diagnosis and treatment, congestive heart failure is the only major cardiovascular disorder which is increasing in incidence. Ventricular arrhythmias account for approximately 50 % of the mortalities associated with congestive heart failure. Ventricular arrhythmias typically arise from prolongation of the action potential duration (APD) which results in unstable repolarization and thus generation of arrhythmias. Treatment in these cases can be effected by shortening the action potential or by synchronizing repolarization. This can be achieved by transplanting cells having potassium channels (e.g. delayed rectifier or ether-go-go) which would function in shortening the cardiomyocytic APD.

Heart failure can also be treated by transplantation of cells having L-type or T-type calcium channels into the ventricles in a diffuse or a predetermined pattern in order to increase the excitability of the ventricles and to modulate calcium ion kinetics in the host myocardial tissue. Such transplantation would improve the contractility and relaxation pattern of the ventricles and thus change the systolic and diastolic properties of the ventricle.

Long QT syndrome: patients with genetic or acquired abnormalities in repolarization which display prolonged QT intervals may suffer from

life-threatening malignant arrhythmias such as polymorphic VT. Such patients may be treated with the cellular grafts of the present invention having ion channels, such as potassium channels, which are selected capable of shortening and homogenizing repolarization.

5

EXAMPLE 2

Pancreas

Diabetes Mellitus is a metabolic disease in which carbohydrate utilization is reduced while utilization of lipid and protein enhanced. Diabetes Mellitus is caused by relative deficiency of insulin, and is characterized, in more severe cases, by chronic hyperglycemia, glycosuria, water and electrolyte loss, and various organ damage causing significant morbidity and mortality.

Gap junctions and junction-mediated cell-to-cell communications are obligatory features of gland cells, regardless of their secretory products. Studies on pancreatic islets and acinar cells indicate that cell-to-cell communication via gap junction channels is required for proper biosynthesis, storage and release of both insulin and amylase. However, the endocrine and exocrine portions of the pancreas show opposite connexin (Cx) and coupling changes in relation to the activation and inhibition of their secretory functions. These differences may be accounted for by the expression of connexin43 (Cx43) in pancreatic islets and of Cx26 and Cx32 in pancreatic acini. This alternative expression of

connexin isoforms is also found in several other endocrine and exocrine glands. These observations indicate that connexin-made channels play a central role in the control of secretory events (Meda, 1996, *Clinical & Experimental Pharmacology & Physiology*, Dec;23(12):1053-7).

5 The function of T-type voltage-gated calcium channels in insulin-secreting cells has been previously described (Bhattacharjee et al, 1997 *Endocrinology*, Sep.138(9):3735-40). Whole-cell voltage and current recordings, capacitance measurements, and RIA techniques were used to determine the contribution of T-type calcium channels in modulation of
10 electrical activity and in stimulus-secretion coupling in a rat insulin secreting cell line, INS-1. Studies employing double pulse protocols in the current-clamp mode, uncovered that activation of T-type calcium channels provided a low threshold depolarizing potential that decreased the latency of onset of action potentials and increased the frequency of action
15 potentials, both of which are abolished by administration of nickel chloride (NiCl_2), a selective T-type calcium channel blocker (Bhattacharjee et al, 1997 *Endocrinology*, Sep.138(9):3735-40).

Currently, treatment of non insulin dependent diabetes mellitus (NIDDM) includes, in more severe cases, drug therapy and insulin
20 injections. The sulfonylureas family acts as ATP-sensitive potassium channels blockers, thus causing depolarization of the pancreatic β cells, calcium influx and insulin secretion.

Cellular grafts capable of forming gap junction (e.g. expressing Cx43) with pancreatic beta cells can be used by the present invention to treat NIDDM. These cells which can be of autogeneic, allogeneic or xenogeneic origin can be, for example, transfected ex-vivo with nucleic acid construct encoding a specific ion channel polypeptide(s), such as, for example, CACNA1G (encoded by GenBank Accession number AF134986) which forms a T-type voltage gated calcium channel (see Table 2 below for additional examples). The cells will be transplanted in the pancreas in a diffuse or a predetermined pattern via invasive or minimally invasive techniques. For example, minimally invasive percutaneous procedures using image guiding (CT, US etc) can be used for transplantation of the cellular grafts.

Upon gap junction establishment, the cellular grafts will form a single compartment with the surrounding tissue and will increase the sensitivity of the pancreatic b cells to glucose levels by increasing the depolarization process and the sensitivity of insulin secretion to depolarization. For example, by using cells transfected with the T-type voltage gated calcium channels one may increase the ca influx following depolarization of the pancreatic cells thereby increasing insulin secretion.

Pharmacological blockage of these channels at a fine tuned dosage will prevent spontaneous action potentials thus preventing hypoglycemic

states. This approach is advantageous since it allows to monitor insulin secretion regardless of the time of drug administration.

Several approaches can be utilized for regulating pancreatic beta cells excitability and insulin secretion. For example, transplantation of cells transfected with sodium or calcium channels can be utilized to increase depolarization of the beta cells or transplantation of cells transfected with calcium channels can be utilized to increase calcium influx thereby increasing beta cell sensitivity to depolarization. in addition these and other approaches can be utilized to increase and prolong the firing rate of such pancreatic cells.

Table 2

Ion	Channel type	GenBank Accession numbers
Na	Sodium channel	AB027567
Na	Voltage gated "SCN11A"	AF188679
Na	hBNC2"; product: "sodium channel 2	U78181 AL035862 AA442069 AI017398 AI620655 AI762424 Z40887 AI700050
Ca	T-type	AF134986
Ca	'VOLTAGE-DEPENDENT N-TYPE CALCIUM CHANNEL'	M94172 U76666 AA776162 T12610
Ca	"L-type calcium channel (HFCC)"; Human'	M92269 AA927640 AA443875 AA173146
Ca	"CACNG4"; product: "calcium channel'	AF142625
Ca	'VOLTAGE-DEPENDENT L-TYPE CALCIUM CHANNEL,'	AJ224874 AJ006216
Ca	"voltage-dependent calcium channel'	M92301 W07059 T28094
Ca	L-type	M76558 AF055575 H29339 R25307 T27949 AA885750 AW029633 AI955764 AW008794 AA978315 AI914244 AI951788 AW008769 H29256 AI963788 AI537488 AA468565 AA523647 AI361691 R46658 AW139850 AI017959 AA701888 AA703120 AA877582

EXAMPLE 3**CNS****Epilepsy:**

5 Epilepsy is a chronic disorder usually associated with some alteration of consciousness and characterized by paroxysmal brain dysfunction due to excessive neuronal discharge.

Astroglial cells contribute to neuronal maintenance and function in the normal and diseased brain. Gap junctions, formed predominantly by
 10 connexin43 between astroglia, provide important pathways which coordinate astroglial responses (Reuss et al, 2000, Glia May;30(3):231-41). Neuronal-glia interactions play an important role in information processing in the CNS. Previous studies have indicated that electro-ionic coupling between locus ceruleus (LC) neurons is involved in
 15 synchronizing the spontaneous activity. Moreover, Spontaneous oscillations in the membrane potential were observed in a subset of glial cells. These oscillations were synchronous with the firing of neurons, insensitive to transmitter receptor antagonists and disrupted by carbenoxolone, a gap junction blocker. Finally, immunoelectron
 20 microscopy studies established that connexins, the proteins that form gap junctions, were present on portions of the plasmalemma, bridging the

cytoplasm of neurons and glia in LC (Alvarez et al, 2000, J Neurosci. Jun 1;20(11):4091-8).

Treatment of epilepsy can be effected by the present invention by transplantation of astroglial cells, fibroblasts or other cells transfected ex-
 5 vivo with a restraining force channel coding sequence exemplified in Table 3 below.

Table 3

Ion	Channel type	GenBank Accession numbers
K	Kv1.3	H18261
K	inward rectifier potassium channel TWIK-1 - human	S65566
K	Delayed rectifier potassium channel - human	L28168 L33815 M26685
K	Cardiac inward rectifier potassium channel - human	I38727
K	VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KV1.4	M55514 A1631014 A1701825 A1694934 A1793138
K	'voltage-gated potassium channel protein - human'	JC5275
K	OKCNQ2"; potassium channel	AF033348
K	'inwardly rectifying potassium channel, hippocampal	I38521
K	VOLTAGE-GATED POTASSIUM CHANNEL PROTEIN KQT-LIKE 3.KCNQ3.	AF033347 AF071491 AW205596 AW135705 AA019129 AA001392 H86059 H08544 R36327 T78692 A1125802 H08545 R49258

The transfected cells will be transplanted to the pathologic foci
 10 using standard neuro-surgical methods. Upon establishment of gap junction with the surrounding tissue, the cellular grafts form a single compartment which enables the repression of pathological tissue regions via controlled activation of the channels.

EXAMPLE 4

Neuronal networks

Neuronal cells were cultured on multi electrode arrays in efforts to determine electrophysiological function of these cultured cells under various conditions.

Culture techniques:

Cortical neurons were obtained from newborn rats within 24 hours from birth, following standard harvesting procedures (Culturing nerve cells, 2nd edition, Gary Ranker and Kimberly Goslin, 1998). The cortex tissue was digested enzymatically and mechanically dissociated and the neurons were plated directly onto substrate-integrated multi-electrode array (MEA) dishes prepared as described below. The cultures were bathed in MEM which was supplemented with heat-inactivated horse serum (5%), Glutamine (0.5 mM), Glucose (20 mM), and Gentamycin (10 μ g/ml), and maintained in a tissue culture incubator at 37 °C, 5 % CO₂ and 95 % during the recording phases. Half of the medium was exchanged twice a week and the experiments were performed during the third week following plating, thus allowing complete maturation of the neurons (Figures 1a-b).

It is a well known fact that electrical activity in a cultured neuronal network is dependent upon synaptic transmission. As shown by various published studies, this electrical activity can be blocked by perfusion with

the *N*-Methyl-D-aspartate (NMDA), receptor antagonist D-2-amino-5-phosphonovalerate (APV), and non-NMDA receptor antagonist 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX).

To determine the sensitivity and accuracy of the multi electrode array and detecting system of the present invention, prior art electrical activity studies in cultured neuronal networks were repeated as part of the present study using intracellular recordings as well as MEA recordings.

Electrophysiological methods:

Arrays of 60 Ti/Au/TiN electrodes, 30 μ m in diameter, spaced 200 μ m from each other (MultiChannelSystems (MCS), Reutlingen, Germany) were utilized in the present study. The insulation layer (silicon nitride), was pretreated with poly-L-lysine forming a good surface for network development. A commercial 60-channel amplifier (B-MEA-1060, MCS, Reutlingen, Germany) with frequency limit range of 10-3000 Hz and a gain of $\times 1024$ was utilized for signal amplification. The amplifier was connected to MCPPlus filter amplifiers (Alpha Omega, Nazareth, Israel) for further amplification ($\times 10$ to $\times 20$). Stimulation through the MEA was performed using a dedicated 8-channel stimulus generator (MCS, Reutlingen, Germany).

In addition, the micro-incubation environment was arranged to support long-term recordings from MEA dishes. This was achieved by streaming a filtered, heated and humidified air/CO₂ (95/5%) gas mixture,

and by electrically heating the MEA platform to 37°C. Data is digitized using two 5200a/526 A/D boards (Microstar Laboratories, WA, USA).

Experiments were first conducted in efforts to determine the functionality of the multi electrode array and the detecting system
5 described above. The response of the cultured neuronal network to electrical stimuli is illustrated in Figures 1c-f.

Following electrical functionality determination, the neuronal network cultures were incubated with various electrical conduction blockers.

10 The addition of 5 μ M bicuculin, 10 μ M DNQX or 20 μ M APV to the cultured neuronal network completely abolished spiking activity therein.

Epilepsy:

Epileptic activity of the cultured neuronal network described above was measured from the MEA described above. Figures 2a-b illustrate
15 epileptic activity recorded from MEA in a mature (3 weeks in vitro) cultured cortical network. This recorded spontaneously bursting synchronous activity throughout the network is a characteristic feature of epileptic-like activity in networks of neurons.

EXAMPLE 5

20 Although electrical coupling between fibroblasts and myocytes has been previously reported by Rook et al. (1992), the experiments conducted as a part of that study were designed in efforts to elucidate the validity of

While reducing the present invention to practice, the present inventors utilized a cell culture model system which included fibroblasts which were transfected with ion channel coding sequences and co-cultured with cardiomyocytes. These co-cultures enabled to test the effects of the ion channel expressing fibroblast on the electrophysiological function of the myocardial cells and to test the effects of various molecules which regulate channel permeability.

Preparation of cultured cardiomyocytes:

Preparation of fibroblast cultures transfected with Kv1.3:

20 Fibroblasts from the NIH 3T3 cell line were transfected with an
expression cassette which included a mutant voltage gated potassium
channel (Kv1.3) coding sequence (GeneBank Accession number H18261)

placed under the transcription control of a constitutive promoter using standard procedures. Fibroblast cultures not transfected with the channel coding sequence were produced from the NIH 3T3 cell-line.

Preparation of co-cultures:

5 Once well synchronous spontaneous activity was established in the cardiomyocyte cultures, fibroblasts transfected with the Kv1.3 channel coding sequence or non-transfected fibroblasts were added to the cultures. Two different methods were used to seed the fibroblasts. In the first method, the fibroblasts were suspended in trypsin for 5 minutes
10 following which they were seeded in a diffuse pattern in the cardiomyocytic cultures (Figure 4a). In the second method, the fibroblasts were pipetted up and down through a 5 ml pipette for 2 minutes and seeded in the cardiomyocytic cultures thus were forming clusters of fibroblasts (Figure 4b).

Immunohistochemistry:

15 The fibroblasts were labeled with Fast DiO [3,3'-dilinoleyloxacarbocyanine perchlorate (*FAST* DiO™ solid, Cat #-3898; Molecular Probes, USA) in order to track the fibroblasts in the co-cultures.

The data acquisition system and electrical activity recording:

20 Extracellular recordings from cultured cardiomyocytes were performed on a PC-based Microelectrode Data Acquisition System (Multi Channel Systems, Reutlingen, Germany), consisting of Multi-Electrode

Array (MEA), pre- and filter-amplifiers, data acquisition board, and software. The MEA consists of a 50×50 mm glass substrate, in the center of which is an embedded 0.7×0.7 or 1.4×1.4 mm matrix of 60 Titanium-nitride, gold contacts 10 or 30 µm diameter electrodes insulated with
 5 silicone nitride, with inter-electrode distance of 100 or 200 µm (there are no electrodes in the corners of the matrix). Data were recorded at 10-25 KHz with 12-bit precision. During the recording sessions, the MEA (removed from the regular incubator) was constantly perfused with a gas mixture consisting of 5% CO₂ + 95% air. Temperature was kept at $37 \pm 0.10^{\circ}\text{C}$.

10 ***Construction of activation maps:***

Recorded data was filtered using cutoff frequency of 2 KHz (Fast et al, 1993). The filtered signal was then differentiated digitally to determine the Local Activation Time (LAT) at each electrode, corresponding to dF/dt_{min} (where F is the filtered signal) (Dolber and Spach, 1986). Color-
 15 coded activation maps were constructed by interpolating the LAT values for the sites between the electrodes, and by extrapolating the LAT values for the 4 corners of the MEA matrix. Activation maps were plotted by means of Matlab standard 2-d plotting function (pcolor) (Matlab 5.3 Mathworks Incorporated©). Conduction velocity was calculated by
 20 standard methods (Bayly et al, 1988).

Conduction block:

Conduction block quantification is central to evaluating conduction block development in the cell cultures, and to evaluating reversibility of the conduction block following CTX application. A conduction block was determined using the following algorithm:

the local activation time (LAT) of each electrode was compared to the LAT of the four nearest electrodes, where LAT_x is the local activation time at electrode x and LAT_y is the local activation time at one of the four nearest electrodes to electrode x. Thus, If $LAT_x - LAT_y > 0.25 \times [LAT_{max}(last$

local activation time in the array)-LAT_{min}(first local activation time in the array)], then the electrode was assigned a value of 1, else the electrode was assigned a value of 0. Each of the four electrodes was tested and if one or more satisfied this condition, then electrode x was set to a value – 1, a sum of all the electrode values represented the block value.

Recording protocol:

Electrical activity of the cultures was recorded on day one immediately prior to seeding of the fibroblasts and then daily until the cultures died or no spontaneous activity was detected. During the daily measurements the cultures were subjected to increasing concentrations

(0.1, 1, 10, 100 nM) of CTX.

RESULTS

Spontaneous activity:

Measurement were performed during spontaneous activity from three groups of cardiomyocyte cultures: cultures without fibroblasts, 5 cultures with NIH 3T3 fibroblasts (seeded diffusely) and cultures with transfected NIH 3T3 fibroblasts (expressing the mutant voltage gated potassium channel coding sequence).

The cultures where subjected to an increasing concentration of CTX from 0.1 to 100 nM. CTX caused a significant increase in the 10 spontaneous activity rate in the co-cultures which included the transfected fibroblasts. The activity rate increased by 20 and 55% following administration of CTX concentrations of 10 and 100 nM respectively. In contrast, administration of CTX to cardiomyocyte cultures or to cardiomyocytes co-cultured with untransfected fibroblasts 15 did not increase the activity rate at 10 nM and caused a modest increase of up to 15% at a 100 nM (Figure 5).

There are three possible explanations for these results:

(i) Kv1.3 channel opening in transfected fibroblasts during action potential propagation causes hyperpolarization and therefore elongation 20 of phase 4 at neighboring cardiomyocytes resulting in a slower activity rate; therefore, blocking of Kv1.3 channels with CTX reverses this effect.

(ii) CTX treatment increases electrical activity in areas that are

blocked due to the presence of fibroblasts.

(iii) CTX enables propagation through otherwise blocked conduction tracts thus enabling propagation of action potentials. Since the area of the electrode array is a $1 \times 1 \text{ mm}^2$ and since the area of the plate in which the array is embedded is about 2 cm^2 , most of the culture activity is not recorded because propagation of electrical signal from cells positioned outside the array may be blocked prior to entering the array.

Application of CTX opens conduction blocks and thus enables activation of the myocytes at the electrode area.

The weak response observed in the control cultures treated with a high concentration of CTX is probably due to a minor blockage of potassium channels in the myocytes.

Synchronous activity and conduction block:

The two control culture types (with or without untransfected fibroblasts) exhibited a well-coupled synchronous activity throughout the experiment. Four of the co-cultures with transfected fibroblasts demonstrated an uncoupling effect following the fibroblasts seeding due to a conduction block generated by the Kv1.3 channels formed in the fibroblasts. Uncoupling effect was reversed following treatment with CTX (Figures 6f-g). Almost all of the cultures which included transfected fibroblasts demonstrated conduction blocks which developed following fibroblasts seeding (Figure 10). Such conduction blocks were reversed

following treatment with CTX (Figure 7d and Figure 11).

Cultures including non-transfected fibroblasts did not demonstrate conduction blocks or reversibility of blocks following application of CTX (Figures 12a-d). An immediate decrease in conduction velocity following transfected fibroblast seeding was also observed. Such an effect was not observed in co-cultures that included non-transfected fibroblasts (Figure 8).

Amplitude change:

In comparison to cultures seeded with non-transfected fibroblasts, the amplitude of the extracellular signals decreased significantly following seeding with transfected fibroblasts (Figure 9). This result may indicate a general decrease in culture excitability, implicating a reduced mass of action potential generating cardiomyocytes or the presence of slow conduction.

Summary and future directions

The above described results demonstrate for the first time that transplantation of fibroblasts transfected with a Kv1.3 channel coding sequence into cardiomyocytic cultures causes a significant change in the electrophysiological function of this excitable tissue.

Specifically, reduced spontaneous rate of the co-culture's excitability, lower amplitude of extracellular potentials, reduced conduction velocity and generations of local conduction blocks, were generated.

These changes were partially or fully reversed following administration of a specific Kv1.3 channel blocker, CTX. These results indicate the presence of tight structural and functional coupling between the fibroblasts and the myocytes, activation of the Kv1.3 channels and significant modulation of the electrical properties of the cultures.

Thus, the present invention provides a novel method which can be utilized to modulate the electrophysiological function of an excitable tissue region, which method can be utilized to treat various cardiac disorders.

The ability to modulate the electrophysiological properties of cardiac tissue may have significant clinical applications. Transplantation of cellular grafts having a predetermined electrical phenotype may be used, in the future, to alter the electrophysiological properties of cardiac tissue and together with pharmacological administration serve as a procedure for treating selected pathologies in the heart.

Furthermore, the method of the present invention is advantageous in that it effects a local tissue region rather than the heart as a whole, thus not affecting non-pathological tissue regions. This mode of treatment may be applied to treat a variety of cardiac arrhythmias.

For examples, transplantation of cellular grafts of the present invention to the AV node may be used for AV nodal modification, where the inherent properties of the cellular graft (the frequency response with or without specific pharmacology) can be used to modify the ventricular

response during different atrial arrhythmias thus replacing the need for pharmacological treatment.

Local transplantation of the cellular grafts of the present invention may also be used to repress arrhythmogenic foci arising due to abnormal automaticity or to repress triggered activity by modulating the action potential in selected tissue regions.

In addition, reentrant arrhythmia may also benefit from the teachings of the present invention. Cellular grafts may be used to create a local conduction block in a critical area of the circuit thus treating the arrhythmia. Alternatively, predetermined seeding patterns may be used to create barriers or lines of conduction blocks for the treatment of more complex reentrant arrhythmias such as atrial fibrillation or flutter. In the later two cases, specific cell types can be used to allow normal conduction during slow (normal) rhythms, while creating local conduction blocks during fast (pathological) rhythms.

The teachings of the present invention may also be applied to modify the electrophysiological functionality of excitable tissues such as, for example, nervous tissue and glandular tissue. For examples, astrocyte transfected with selected ion channels may be used to modulate focal pathological areas in the CNS, thus enabling treatment of disorders such as epilepsy Parkinson and the like.

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims. All publications, patents, patent applications and sequences disclosed therein and/or identified by a GeneBank accession number mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent, patent application or sequence was specifically and individually indicated to be incorporated herein by reference. In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention.

REFERENCES

1. Bayly P.V, Bruce H.Ken Knight, Jack M. Rogers, Russel E. Hillsley, Raymond E. Ideker, William M. Smith. Estimation of conduction velocity vector fields from epicardial mapping data. IEEE transactions on biomedical engineering. 1988. 45:563-571.
2. Fast VG, Kleber AG. Microscopic conduction in cultured strands of neonatal rat heart cells measured with voltage-sensitive dyes. Circ Res. 1993. 73: 914-925.
3. Gussoni, E., Pavlath, G. K., Lanctot, A. M., Sharma, K. R., Miller, R. G., Steinman, L. and Blau, H. M. Normal dystrophin transcripts detected in duchenne muscular dystrophy patients after myoblast transplantation. Nature. 1992. 356: 435-438.
4. Marom s, Goldstein SA, Kupper J, Levitan IB. Mechanism and modulation of inactivation of the Kv3 potassium channel. *Receptor and Channels*. 1993. 1:81-88.
5. Rook MB, Van Ginneken ACG, De Jonge B, Aoumari AE, Gros D, and Jongsma HJ. Differences in gap junction channels between cardiac myocytes, fibroblasts, and heterologous pairs. Am J Physiol. 1992. 263: C959-C977.
6. Rubin Y, Kessler-Icekson G, Navon G. The effect of furosemide on calcium ion concentration in myocardial cells. Cell Calcium. 1995 Aug;18(2):135-9.

7. Spach MS, Dolber PC. Relating extracellular potentials and their derivatives to anisotropic propagation at a microscopic level in human cardiac muscle. Evidence for electrical uncoupling of side-to-side fiber connections with increasing age. *Circ Res.* 1986 Mar;58(3):356-71.
8. Tompson, L. Fetal transplants show promise. *Science.* 1992. 257: 868-870.

000207" 68876960

WHAT IS CLAIMED IS:

1. A nucleic acid construct comprising:
 - (a) a first polynucleotide region encoding at least one first polypeptide capable of forming a functional ion channel or transporter when expressed within a cell; and
 - (b) a second polynucleotide region encoding at least one second polypeptide capable of forming a functional gap junction when expressed within said cell.
2. A cell, cell culture or tissue explant transformed with the nucleic acid construct of claim 1.
3. The cell, cell culture or tissue explant of claim 2, wherein the cell is selected from the group consisting of a fibroblast, a myoblast, an astroglial cell and an endothelial cell.
4. The cell, cell culture or tissue explant of claim 2, wherein the tissue explant is an organ tissue explant.

5. A pharmaceutical composition comprising, as an active ingredient, the nucleic acid construct of claim 1 and a pharmaceutically acceptable carrier.

6. The nucleic acid construct of claim 1, wherein said ion channel is selected from the group consisting of a sodium ion channel, a potassium ion channel, a calcium ion channel and a chloride ion channel.

7. The nucleic acid construct of claim 1, wherein said at least one first polypeptide is selected from the group consisting of a K channel polypeptide, a Na channel polypeptide, a Ca channel polypeptide, a Cl channel polypeptide, a Na/K transporter polypeptide, a Na/Ca transporter polypeptide, a Na/H transporter polypeptide and a Cl/HCO₃ transporter polypeptide

8. The nucleic acid construct of claim 1, wherein said at least one second polypeptide is selected from the group consisting of connexin43, connexin45 and connexin26.

9. The nucleic acid construct of claim 1, further comprising at least one promoter being for directing the transcription of said first polynucleotide and said second polynucleotide.

10. The nucleic acid construct of claim 9, wherein said at least one promoter is functional in mammalian cells.

11. The nucleic acid construct of claim 9, wherein said at least one promoter is selected from the group consisting of a constitutive promoter, a tissue specific promoter, an inducible promoter and a developmentally regulated promoter.

12. The nucleic acid construct of claim 1, wherein said first polynucleotide region and said second polynucleotide region are transcriptionally fused via an IRES sequence.

13. The nucleic acid construct of claim 1, wherein said at least one first polypeptide and said at least one second polypeptide are translationally fused via at least one protease recognition site.

14. The nucleic acid construct of claim 9, wherein said at least one promoter includes two promoters, a first promoter for directing the transcription of said first polynucleotide and a second promoter for directing the transcription of said second polynucleotide.

15. A nucleic acid construct system comprising:
- (a) a first nucleic acid construct including a first polynucleotide region encoding at least one first polypeptide capable of forming a functional ion channel or transporter when expressed within a cell; and
 - (b) a second nucleic acid construct including a second polynucleotide region encoding at least one second polypeptide capable of forming a functional gap junction when expressed within said cell.
16. A cell, cell culture or tissue explant transformed with the nucleic acid construct of claim 15.
17. A pharmaceutical composition comprising, as an active ingredient, the nucleic acid construct of claim 15 and a pharmaceutically acceptable carrier.
18. The nucleic acid construct system of claim 15, wherein said first nucleic acid construct further includes a first promoter being for directing the transcription of said first polynucleotide and further wherein

000201 68816960

said second nucleic acid construct further includes a second promoter being for directing the transcription of said second polynucleotide.

19. The nucleic acid construct system of claim 18, wherein each of said first and said second promoters is functional in mammalian cells.

20. The nucleic acid construct system of claim 18, wherein each of first and said second promoters is independently selected from the group consisting of a constitutive promoter, a tissue specific promoter, an inducible promoter and a developmentally regulated promoter.

21. The nucleic acid construct system of claim 15, wherein said at least one first polypeptide is selected from the group consisting of a K channel polypeptide, a Na channel polypeptide, a Ca channel polypeptide, a Cl channel polypeptide, a Na/K transporter polypeptide, a Na/Ca transporter polypeptide, a Na/H transporter polypeptide and a Cl/ HCO₃ transporter polypeptide.

22. The nucleic acid construct system of claim 15, wherein said at least one second polypeptide is selected from the group consisting of connexin43, connexin45 and connexin 26.

23. A method of modifying the electrophysiological function of an excitable tissue region of an individual, the method comprising the step of implanting cells into the excitable tissue region, each implanted cell being:

- (a) capable of forming gap junctions with at least one cell of the excitable tissue region; and
- (b) capable of forming a functional ion channel or transporter said functional ion channel or transporter being capable of modifying the electrophysiological function of the excitable tissue region.

24. The method of claim 23, wherein said ion channel is selected from the group consisting of a sodium ion channel, a potassium ion channel, a calcium ion channel and chloride ion channel.

25. The method of claim 23, wherein each implanted cell is transformed, prior to, or following implantation, with an exogenous polynucleotide expressing at least one polypeptide capable of forming said functional ion channel or transporter.

26. The method of claim 23, wherein each implanted cell is transformed, prior to, or following implantation, with an exogenous

polynucleotide expressing at least one polypeptide capable of forming said gap junctions.

27. The method of claim 25, wherein expression of said at least one polypeptide from said exogenous polynucleotide is regulatable by an endogenous or an exogenous factor.

28. The method of claim 23, wherein an ion permeability of said functional ion channel is regulatable by an endogenous or an exogenous factor.

29. The method of claim 23, further comprising the step of regulating permeability of said functional ion channel or an activity of said transporter to thereby regulate the electrophysiological function of the excitable tissue region.

30. The method of claim 28, wherein said step of regulating said permeability is effected by administering said exogenous factor to the excitable tissue region.

31. The method of claim 23, wherein each implanted cell is capable of forming said functional ion channel or transporter following induction.

32. The method of claim 23, wherein the excitable tissue region forms a part of an organ selected from the group consisting of a heart, a pancreas, a kidney, a brain and a liver.

33. The method of claim 23, wherein the method is utilized for regulating cardiac arrhythmia.

34. The method of claim 23, wherein the method is utilized for regulating secretion of endogenous factors from an organ including the excitable tissue region of the individual.

35. The method of claim 23, wherein the method is utilized for regulating neuronal discharge.

36. A method of modifying the electrophysiological function of an excitable tissue region of an individual, the method comprising the step of expressing an exogenous polypeptide in at least a portion of cells forming a part of, or being in contact with, the excitable tissue region, said

10

37. The method of claim 36, further comprising the step of expressing a second exogenous polypeptide in said at least a portion of said cells, said second exogenous polypeptide being capable of forming functional gap junctions within said at least a portion of said cells.

The following year	
1990	1991
1992	1993
1994	1995
1996	1997
1998	1999
2000	2001
2002	2003
2004	2005
2006	2007
2008	2009
2010	2011
2012	2013
2014	2015
2016	2017
2018	2019
2020	2021
2022	2023
2024	2025
2026	2027
2028	2029
2030	2031
2032	2033
2034	2035
2036	2037
2038	2039
2040	2041
2042	2043
2044	2045
2046	2047
2048	2049
2050	2051
2052	2053
2054	2055
2056	2057
2058	2059
2060	2061
2062	2063
2064	2065
2066	2067
2068	2069
2070	2071
2072	2073
2074	2075
2076	2077
2078	2079
2080	2081
2082	2083
2084	2085
2086	2087
2088	2089
2090	2091
2092	2093
2094	2095
2096	2097
2098	2099
2100	2101
2102	2103
2104	2105
2106	2107
2108	2109
2110	2111
2112	2113
2114	2115
2116	2117
2118	2119
2120	2121
2122	2123
2124	2125
2126	2127
2128	2129
2130	2131
2132	2133
2134	2135
2136	2137
2138	2139
2140	2141
2142	2143
2144	2145
2146	2147
2148	2149
2150	2151
2152	2153
2154	2155
2156	2157
2158	2159
2160	2161
2162	2163
2164	2165
2166	2167
2168	2169
2170	2171
2172	2173
2174	2175
2176	2177
2178	2179
2180	2181
2182	2183
2184	2185
2186	2187
2188	2189
2190	2191
2192	2193
2194	2195
2196	2197
2198	2199
2200	2201
2202	2203
2204	2205
2206	2207
2208	2209
2210	2211
2212	2213
2214	2215
2216	2217
2218	2219
2220	2221
2222	2223
2224	2225
2226	2227
2228	2229
2230	2231
2232	2233
2234	2235
2236	2237
2238	2239
2240	2241
2242	2243
2244	2245
2246	2247
2248	2249
2250	2251
2252	2253
2254	2255
2256	2257
2258	2259
2260	2261
2262	2263
2264	2265
2266	2267
2268	2269
2270	2271
2272	2273
2274	2275
2276	2277
2278	2279
2280	2281
2282	2283
2284	2285
2286	2287
2288	2289
2290	

A method of modifying the electrophysiological function of an excitable tissue region of an individual is provided. The method includes the step of implanting cells into the excitable tissue region. Each implanted cell is (a) capable of forming gap junctions with at least one cell of the excitable tissue region; and (b) capable of forming a functional ion channel or transporter, wherein the functional ion channel or transporter is capable of modifying the electrophysiological function of the excitable tissue region.

Fig 1a




Fig 1b

Fig 1c




Fig 1d




Fig 1e




Fig 1f




Fig 2a




Fig 2b





Fig. 3



Fig 4a



Fig 4b






Fig 5



Figs 6a-g

000207 6806060



Figs 7a-d




Fig 8




Fig 9

000207 68876960

conduction
block




Fig 10

conduction
block






Fig 11

000207" 688F6960


12A




12B



12C



12D



Figs. 12a-d

Docket No.
00/20989**Declaration and Power of Attorney For Patent Application****English Language Declaration**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

**NUCLEIC ACID CONSTRUCTS AND CELLS, AND METHOD UTILIZING SAME
FOR MODIFYING THE ELECTROPHYSIOLOGICAL FUNCTION OF EXCITABLE
TISSUES**

the specification of which

☒ is attached hereto.

☐ was filed on _____ as United States Application No. or PCT

International Application Number _____

and was amended on _____

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

(Number)

(Country)

(Day/Month/Year Filed)

☐

(Number)

(Country)

(Day/Month/Year Filed)

☐

(Number)

(Country)

(Day/Month/Year Filed)

☐

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

(Application Serial No.)

(Filing Date)

(Application Serial No.)

(Filing Date)

(Application Serial No.)

(Filing Date)

I hereby claim the benefit under 35 U.S.C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112. I acknowledge the duty to disclose to the United States Patent and Trademark Office all the information known to me to be material to patentability as defined in Title 37, C.F.R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*

SOL SHEINBEIN Registration Number 25,457

MARTIN MOYNIHAN Registration Number 40,338

Send Correspondence to:

**G.E. EHRLICH (1995) LTD.
c/o ANTHONY CASTORINA
SUITE 207
2001 JEFFERSON DAVIS HIGHWAY
ARLINGTON, VIRGINIA 22202, USA**

Direct Telephone Calls to: *(name and telephone number)*

Anthony Castorina

Tel. No. 703-4151581

FULL NAME OF SOLE OR FIRST INVENTOR

YAIR FELD

Sole or first inventor's signature

Date 16 October 2000

Residence : 10 Hasport, Haifa, Israel

Citizenship : ISRAELI

Post Office Address : 10 Hasport, Haifa, Israel

FULL NAME OF SECOND INVENTOR, IF ANY,

LIOR GEPSTEIN

Second inventor's signature

Date 16 October 2000

Residence : 26 Raanan Street, Haifa, Israel

Citizenship : ISRAELI

Post Office Address : 26 Raanan Street, Haifa, Israel

FULL NAME OF THIRD INVENTOR , IF ANY		SHIMON MAROM
Third inventor's signature <i>שמואל מרומ</i>		Date <u>16 October 2000</u>
Residence	:	6 Ovadya Street, Haifa, Israel
Citizenship	:	ISRAELI
Post Office Address	:	6 Ovadya Street, Haifa, Israel

FULL NAME OF FOURTH INVENTOR , IF ANY		MEIRA FRANK
Fourth inventor's signature <i>מירה פרנק</i>		Date <u>16 October 2000</u>
Residence	:	3 Yanush Kortchack Street, Haifa, Israel
Citizenship	:	ISRAELI
Post Office Address	:	3 Yanush Kortchack Street, Haifa, Israel